

AD-A186 387


ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE  
EIGENVALUES OF RANDOM MATRIC (U) PITTSBURGH UNIV PA  
CENTER FOR MULTIVARIATE ANALYSIS Z D BAI ET AL JUN 87

171

UNCLASSIFIED

TR-87-16 AFOSR-TR-87-1076 F49620-85-C-0008 F/G 12/2

NL





AD-A186 387

DTIC FILE COPY

AFOSR-TR. 87-1076


ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE  
EIGENVALUES OF RANDOM MATRICES WHICH ARISE  
UNDER COMPONENTS OF COVARIANCE MODEL

Z. D. Bai, P. R. Krishnaiah and L. C. Zhao

Center for Multivariate Analysis  
University of Pittsburgh

**Center for Multivariate Analysis**

**University of Pittsburgh**



DTIC  
ELECTE  
SEP 29 1987  
S D  
E

This document has been approved  
for public release and sale by  
Distribution & Marketing

87 9 25 00

①

ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE  
EIGENVALUES OF RANDOM MATRICES WHICH ARISE  
UNDER COMPONENTS OF COVARIANCE MODEL


Z. D. Bai, P. R. Krishnaiah and L. C. Zhao

Center for Multivariate Analysis  
University of Pittsburgh

June 1987

Technical Report No. 87-16

Center for Multivariate Analysis  
Fifth Floor Thackeray Hall  
University of Pittsburgh  
Pittsburgh, PA 15260



Research sponsored by the Air Force Office of Scientific Research (AFSC).  
under Contract F49620-85-C-0008. The United States Government is  
authorized to reproduce and distribute reprints for governmental purposes  
notwithstanding any copyright notation hereon.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE EIGENVALUES  
OF RANDOM MATRICES WHICH ARISE UNDER COMPONENTS  
OF COVARIANCE MODEL

Z. D. Bai, P. R. Krishnaiah and L. C. Zhao

ABSTRACT

In this paper, the authors derived asymptotic joint distributions of the eigenvalues of some random matrices which arise under components of covariance model.

AMS 1980 subject classifications. Primary 62E20; Secondary 62H15, 62H12.

Key Words and Phrases: Component of covariance model, eigenstructure analysis, limiting distribution, random matrix.

Research sponsored by the Air Force Office of Scientific Research (AFSC).  
under Contract F49620-85-C-0008. The United States Government is  
authorized to reproduce and distribute reprints for governmental purposes  
notwithstanding any copyright notation hereon.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-87-1076</b> <del>87-16</del>	2. GOVT ACCESSION NO. <b>A186 387</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>On the asymptotic joint distributions of the eigenvalues of random matrices which arise under components of covariance model</b>		5. TYPE OF REPORT & PERIOD COVERED <b>Technical - June 1987 Journal</b>
7. AUTHOR(s) <b>Z. D. Bai, P. R. Krishnaiah and L. C. Zhao</b>		6. PERFORMING ORG. REPORT NUMBER <b>87-16</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Center for Multivariate Analysis 515 Thackeray Hall University of Pittsburgh, Pittsburgh, PA 15260</b>		8. CONTRACT OR GRANT NUMBER(s) <b>F49620-85-C-0008</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61021 230445</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>Some C. 11</b>		12. REPORT DATE <b>June 1987</b>
		13. NUMBER OF PAGES <b>16</b>
		15. SECURITY CLASS. (of this report) <b>Unclassified</b>
16. DISTRIBUTION STATEMENT (of this Report) <b>Approved for public release; distribution unlimited</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>Component of covariance model, eigenstructure analysis, limiting distribution, random matrix.</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>In this paper, the authors derived asymptotic joint distributions of the eigenvalues of some random matrices which arise under components of covariance model.</b>		

-A-

ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE EIGENVALUES  
OF RANDOM MATRICES WHICH ARISE UNDER COMPONENTS  
OF COVARIANCE MODEL

Z. D. Bai, P. R. Krishnaiah and L. C. Zhao

ABSTRACT

In this paper, the authors derived asymptotic joint distributions of the eigenvalues of some random matrices which arise under components of covariance model. (Keywords: eigenstructure analysis; multivariate analysis; analysis of variance)

Accession For	
NTIS GRA&I <input type="checkbox"/>	
DTIC TAB <input type="checkbox"/>	
Unannounced <input type="checkbox"/>	
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



AMS 1980 subject classifications. Primary 62E20; Secondary 62H15, 62H12.

Key Words and Phrases: Component of covariance model, eigenstructure analysis, limiting distribution, random matrix.

# ON THE ASYMPTOTIC JOINT DISTRIBUTIONS OF THE EIGENVALUES OF RANDOM MATRICES WHICH ARISE UNDER COMPONENTS OF COVARIANCE MODEL

Z. D. Bai, P. R. Krishnaiah and L. C. Zhao

## 1. INTRODUCTION

Consider the following components of covariance model

$$\underline{x}_{ij} = \underline{\mu} + \underline{\alpha}_i + \underline{\varepsilon}_{ij} \quad (1.1)$$

for  $j=1,2,\dots,m_i$  and  $i=1,2,\dots,k$ , where  $\underline{x}_{ij}: p \times 1$ , is the  $j$ -th observation of the  $i$ -th group,  $\underline{\mu}$  is an unknown general mean vector,  $\underline{\alpha}_i$  is the random effect vector of the  $i$ -th group, and  $\underline{\varepsilon}_{ij}$  is the error vector. It is assumed that  $\underline{\alpha}_i$ 's are iid.,  $\underline{\varepsilon}_{ij}$ 's are iid.,  $\underline{\alpha}_i$ 's and  $\underline{\varepsilon}_{ij}$ 's are independent, and that

$$\begin{aligned} E(\underline{\alpha}_i) &= E(\underline{\varepsilon}_{ij}) = \underline{0}, \\ E(\underline{\alpha}_i \underline{\alpha}_i') &= \underline{\gamma}, \quad E(\underline{\varepsilon}_{ij} \underline{\varepsilon}_{ij}') = \underline{\Sigma}_i > \underline{0}, \\ E(\underline{\alpha}_i \underline{\alpha}_i')^2 + E(\underline{\varepsilon}_{ij} \underline{\varepsilon}_{ij}')^2 &< \infty, \end{aligned} \quad (1.2)$$

where  $\underline{\gamma}$  is a  $p \times p$  non-negative definite matrix. In contrast to the MANOVA model with the non-random effects, which is called the model I, the model (1.1) is also called the model II. If  $m_1 = m_2 = \dots = m_k = m$ , the model is balanced, otherwise unbalanced.

Recently, the components of covariance model has received considerable attention in literature. Its importance not only consists in the generality of the model, but also its connection to linear structural



relationship models and the signal processing. For a discussion on applications in linear structural relations, we refer to Anderson (1984) whereas we refer to Zhao, Krishnaiah and Bai (1986b) for applications in signal processing.

Many papers were devoted to the balanced components of covariance models. Anderson, Anderson and Olkin (1986) studied the maximum likelihood estimators (MLE) of  $\mu$ ,  $\Psi$  and  $\Sigma_1$  under the condition that  $\Psi \geq 0$  and is of a maximum rank, and obtained the likelihood ratio criterion (LRC) for testing the rank hypothesis on  $\Psi$ . Some of these have been obtained by Theobald (1975), and Schott and Saw (1984). Rao (1983) considered some test procedures for testing relationships between two covariance matrices and these procedures are useful in finding the rank of the component of covariance. Some results for the case of unrestricted covariance matrix were announced by Anderson (1984), where an extensive list of references is given. Some authors put the determination of the rank of  $\Psi$  in the framework of model selection; some references of this aspect are given in Zhao, Krishnaiah and Bai (1986 a, 1986 b). In these two papers, they also proposed some procedures to determine the multiplicities of the eigenvalues of the matrix  $\Psi \Sigma_1^{-1}$  or  $\Psi$ .

For the unbalanced components of covariance model, Bai, Krishnaiah and Zhao (1987) proposed some consistent procedures for the determination of the rank  $q$  of the random effect covariance matrix  $\Psi$ . One important case under consideration is that  $\Sigma_1$  is arbitrary definite matrix and  $0 \leq q \leq p$ . It is assumed that

$$1 \leq m_i \leq M, \quad i=1,2,\dots, \quad (1.3)$$

and that

$$k/N \leq \gamma < 1, \quad (1.4)$$

where  $N = \sum_{i=1}^k m_i$ ,  $M$  and  $\gamma$  are constants,  $M$  an integer. We shall keep this assumption in this paper.

Write

$$\bar{X}_i = \sum_{j=1}^{m_i} X_{ij} / m_i, \quad \bar{X}_* = \sum_{i=1}^k m_i \bar{X}_i / N, \quad (1.5)$$

and

$$W_1 = \sum_{i=1}^k \sum_{j=1}^{m_i} (X_{ij} - \bar{X}_i)(X_{ij} - \bar{X}_i)',$$

$$W_2 = \sum_{i=1}^k m_i (\bar{X}_i - \bar{X}_*)(\bar{X}_i - \bar{X}_*)'. \quad (1.6)$$

$W_2$  and  $W_1$  are called the between groups and within group sums of squares and cross products matrices respectively. It is easily seen that

$$\frac{1}{N-k} E(W_1) = \Sigma_1, \quad \frac{1}{k-1} E(W_2) = \Sigma_1 + \frac{N}{k-1} \left(1 - \frac{1}{N^2} \sum_{i=1}^k m_i^2\right) \Psi.$$

We define

$$\Sigma_2 = \Sigma_1 + \frac{N}{k} \Psi, \quad (1.7)$$

which equals  $\frac{1}{k-1} E(W_2)$  when  $m_1 = m_2 = \dots = m_k$ .

In the above literatures, eigenstructure method occupies an important position. Especially, the study for the eigenstructure of the random matrix

$\frac{1}{k-1} W_2 \left( \frac{1}{N-k} W_1 \right)^{-1}$  is of theoretical and practical interest. In this paper, we shall seek for the limiting joint distribution of its eigenvalues.

For the model I, the joint distributions or asymptotic joint distributions of the multivariate analysis of variance (MANOVA) matrices have been found by some authors. Fisher (1939), Hsu (1939) and Roy (1939) have independently derived the joint distribution of the eigenvalues of the MANOVA matrix in the central case. Hsu (1941) derived the above distribution in the noncentral case when the sample size tends to infinity and the underlying distribution is multivariate normal. Bai, Krishnaiah and Liang (1984) extended this result to the case when the underlying distribution is not necessarily multivariate normal. In proving these results, they all assumed that the ratios of the sample sizes of the groups to the total sample size tend to constants in the limiting case, whereas the number of the groups is fixed. On the contrary, in order to seek for the limiting distribution of the eigenvalues of  $\frac{N-k}{k-1}W_2W_1^{-1}$  in the model II, we need to assume that the number  $k$  of the groups tends to infinity.

The balanced and unbalanced cases are discussed in Sections 2 and section 3 respectively.

## 2. THE BALANCED CASE

In this section, we seek for the asymptotic joint distribution of the eigenvalues of the matrix  $\frac{N-k}{k-1} W_2 W_1^{-1}$  for the balanced case. Here we assume that  $k \rightarrow \infty$  and

$$m_1 = m_2 = \dots = m_k = m \geq 2. \quad (2.1)$$

This problem can be reduced to the problem of seeking for the asymptotic joint distribution of the eigenvalues of  $S_2 S_1^{-1}$ , where

$$S_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} \underline{Y}_i \underline{Y}_i', \quad (2.2)$$

$$S_2 = \frac{1}{n} \sum_{i=1}^n \underline{Z}_i \underline{Z}_i', \quad (2.3)$$

and  $\underline{Y}_1, \underline{Y}_2, \dots, \underline{Y}_{n_1}, \underline{Z}_1, \dots, \underline{Z}_n$  are independent  $p \times 1$  vectors,  $\underline{Y}_1, \dots, \underline{Y}_{n_1}$  have a common distribution, so do  $\underline{Z}_1, \dots, \underline{Z}_n$ , satisfying the following conditions:

$$\begin{aligned} E \underline{Y}_1 &= E \underline{Z}_1 = \underline{0}, \\ E \underline{Y}_1 \underline{Y}_1' &= \Sigma_1 > 0, \quad E \underline{Z}_1 \underline{Z}_1' = \Sigma_2 \geq 0, \\ E(\underline{Y}_1' \underline{Y}_1)^2 &< \infty, \quad E(\underline{Z}_1' \underline{Z}_1)^2 < \infty. \end{aligned} \quad (2.4)$$

Without loss of generality, we can assume that  $\Sigma_1 = I_p$  and  $\Sigma_2 = \text{diag} [\lambda_1, \dots, \lambda_p]$  with  $\lambda_1 \geq \dots \geq \lambda_p \geq 0$ , where  $\lambda_1 \geq \dots \geq \lambda_p$  are eigenvalues of  $\Sigma_2 \Sigma_1^{-1}$ .

Write

$$\lambda_i = \varepsilon_h, \quad \text{if } i = a_{h-1} + 1, \dots, a_h, \quad h = 1, 2, \dots, H,$$

where  $\varepsilon_1 > \varepsilon_2 > \dots > \varepsilon_H \geq 0, \quad a_0 = 0, \quad a_h = a_{h-1} + \mu_h,$

$$a_H = p = \mu_1 + \dots + \mu_H.$$

Assume that  $n_1$  varies as  $n$  and  $\lim_{n \rightarrow \infty} n/n_1 = \theta^2$ ,  $0 < \theta^2 < \infty$ . Let

$\phi_1^{(n)} \geq \dots \geq \phi_p^{(n)}$  denote the eigenvalues of  $S_2 S_1^{-1}$ . We are interested in the asymptotic joint distribution of  $\zeta_i^{(n)}$ ,  $i=1,2,\dots,p$ , where

$$\zeta_i^{(n)} = \sqrt{n}(\phi_i^{(n)} - \lambda_i).$$

We need the following lemmas:

LEMMA 2.1. Let  $\underline{X}_n$ ,  $n=0,1,2,\dots$ , be a sequence of random  $p$ -vectors with  $\underline{X}_n \rightarrow \underline{X}_0$  in distribution. Then there exists a probability space  $(\Omega, \mathcal{F}, P)$  on which we can define a sequence of random vectors  $\tilde{\underline{X}}_n$ ,  $n=0,1,2,\dots$ , such that

1.  $\underline{X}_n$  and  $\tilde{\underline{X}}_n$  are identically distributed.
2.  $\tilde{\underline{X}}_n \rightarrow \tilde{\underline{X}}_0$  pointwise.

The above lemma is given in Skorokhod (1956).

LEMMA 2.2. Let  $g_n(x)$  be a sequence of  $K$ -degree polynomials with roots  $x_1^{(n)}, \dots, x_K^{(n)}$  for each  $n$ , and let  $g(x)$  be a  $k$ -degree polynomial with roots  $x_1, \dots, x_k$ ,  $k \leq K$ . If  $g_n(x) \rightarrow g(x)$  as  $n \rightarrow \infty$ , then after suitable rearrangement of  $x_1^{(n)}, \dots, x_K^{(n)}$ , we have  $x_j^{(n)} \rightarrow x_j$ ,  $j=1,\dots,k$  and  $|x_j^{(n)}| \rightarrow \infty$ ,  $j=k+1,\dots,K$ .

For a proof, we refer to Bai (1984).

Put  $n_1 = \theta^{-2} n \beta_n^{-2}$  with  $\beta_n \rightarrow 1$ . The eigenvalues of  $S_2 S_1^{-1}$  are the roots of the determinant equation

$$\det(S_2 - \phi S_1) = 0. \quad (2.5)$$

Write

$$U_n = (u_{ij}^{(n)}) = \sqrt{n_1}(S_1 - \Sigma_1) = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} (Y_i Y_i' - I_p),$$

$$V_n = (v_{ij}^{(n)}) = \sqrt{n}(S_2 - \Sigma_2) = \frac{1}{\sqrt{n}} \sum_{i=1}^n (Z_i Z_i' - \text{diag}[\lambda_1, \dots, \lambda_p]).$$

Then (2.5) becomes

$$\det \left( \frac{1}{\sqrt{n}} V_n - \frac{\theta \beta_n}{\sqrt{n}} \phi U_n - D \right) = 0, \quad (2.6)$$

where

$$D = \begin{pmatrix} (\phi - \xi_1) I_{\mu_1} & & \\ & \ddots & \\ & & (\phi - \xi_H) I_{\mu_H} \end{pmatrix}$$

By the central limit theorem and Lemma 2.1, we can assume that

$$\begin{aligned} U_n &\rightarrow U \triangleq (u_{ij}), \quad \text{a.s.} \\ V_n &\rightarrow V \triangleq (v_{ij}), \quad \text{a.s.} \end{aligned} \quad (2.7)$$

satisfying the following conditions:

- (1)  $\{u_{ij}, 1 \leq i \leq j \leq p\}$  and  $\{v_{ij}, 1 \leq i \leq j \leq p\}$  are two  $p(p+1)/2$  dim. normal vectors independent each other.
- (2)  $u_{ij} = u_{ji}, v_{ij} = v_{ji}$  for all  $i, j$ .
- (3)  $EU_{ij} = Ev_{ij} = 0$  for all  $i, j$ .
- (4)  $\text{cov}(v_{ii}, v_{jj}) = EZ_{1i}^2 Z_{1j}^2 - \lambda_i \lambda_j$ , if  $1 \leq i \leq j \leq p$ ,  
 $\text{cov}(u_{ii}, u_{jj}) = EY_{1i}^2 Y_{1j}^2 - 1$ , if  $1 \leq i \leq j \leq p$ ,

and

$$\text{cov}(v_{i_1 i_2}, v_{j_1 j_2}) = EZ_{1i_1} Z_{1i_2} Z_{1j_1} Z_{1j_2}$$

$$\text{cov}(u_{i_1 i_2}, u_{j_1 j_2}) = EY_{1i_1} Y_{1i_2} Y_{1j_1} Y_{1j_2},$$

if  $1 \leq i_1 \leq i_2 \leq p$ ,  $1 \leq j_1 \leq j_2 \leq p$  and at least one of the strict inequalities  $i_1 < i_2$  and  $j_1 < j_2$  holds. Here  $Y_{1i}$  and  $Z_{1i}$  are the  $i$ -th components of the vectors  $\underline{Y}_1$  and  $\underline{Z}_1$ , respectively.

Split the matrices  $U_n$ ,  $V_n$ ,  $U$  and  $V$  into blocks as follows

$$U_n = (u_{hg}^{(n)}), \quad V_n = (v_{hg}^{(n)}), \quad U = (u_{hg}), \quad V = (v_{hg}),$$

$$h, g = 1, 2, \dots, H,$$

where  $u_{hg}^{(n)}$ ,  $v_{hg}^{(n)}$ ,  $u_{hg}$ ,  $v_{hg}$  have order  $\mu_h \times \mu_g$ ,  $h, g = 1, \dots, H$ .

Take the variable transformation  $\phi = \xi_1 + \tau/\sqrt{n}$ . Multiply by  $n^{1/4}$  the first  $\mu_1$  rows and the first  $\mu_1$  columns of the determinant on the left hand side of (2.6), (2.6) can be rewritten as

$$f_n(\tau) = 0, \quad (2.8)$$

where  $f_n(\tau)$  is a polynomial in  $\tau$  with degree  $p$ , and

$$\lim_{n \rightarrow \infty} f_n(\tau) = f(\tau) =$$

$$= \det \begin{pmatrix} v_{11}^{-\theta \xi_1} u_{11}^{-\tau} I_{\mu_1} & & & \\ & (\xi_2 - \xi_1) I_{\mu_2} & & \\ & & \ddots & \\ & & & (\xi_H - \xi_1) I_{\mu_H} \end{pmatrix}. \quad (2.9)$$

After the variable transformation, the  $p$  roots of (2.6) (i.e., (2.8))

can be written as  $\tau_{i1}^{(n)} = \sqrt{n}(\phi_i^{(n)} - \xi_1)$ . Since  $\lim_{n \rightarrow \infty} \phi_i^{(n)} = \lambda_i$ ,  $i=1, 2, \dots, p$ ,

we have  $\lim_{n \rightarrow \infty} \tau_{i1}^{(n)} = -\infty$ ,  $i=a_1+1, \dots, p$ .

By Lemma 2.2,

$$\lim_{n \rightarrow \infty} \zeta_{i1}^{(n)} = \zeta_i, \quad i=1,2,\dots,a_1, \quad (2.10)$$

where  $\zeta_1, \dots, \zeta_{a_1}$  are the roots of

$$\det (V_{11} - \theta \xi_1 U_{11} - \zeta I_{\mu_1}) = 0. \quad (2.11)$$

Similarly, we can prove that the joint distribution of  $\zeta_1^{(n)}, \dots, \zeta_p^{(n)}$  tends to that of  $\zeta_1, \dots, \zeta_p$  as  $n \rightarrow \infty$ , where  $\zeta_{a_{h-1}+1} \geq \zeta_{a_{h-1}+2} \geq \dots \geq \zeta_{a_h}$  are the roots of

$$\det (V_{hh} - \theta \xi_h U_{hh} - \zeta I_{\mu_h}) = 0, \quad h=1,2,\dots,H \quad (2.12)$$

Thus, we have proved the following

**THEOREM 2.1.** Let  $S_1, S_2$  be defined by (2.2) and (2.3) respectively and let  $\phi_1^{(n)}, \dots, \phi_p^{(n)}$  be the eigenvalues of  $S_2 S_1^{-1}$ . Write  $\zeta_i^{(n)} = \sqrt{n}(\phi_i^{(n)} - \lambda_i)$ ,  $i=1,2,\dots,p$ . Suppose that  $\underline{Y}_1, \dots, \underline{Y}_{n_1}, \underline{Z}_1, \dots, \underline{Z}_n$  are independent and  $\underline{Y}_1, \dots, \underline{Y}_{n_1}$  have a common distribution, so do  $\underline{Z}_1, \dots, \underline{Z}_n$ , where  $E \underline{Y}_1 = E \underline{Z}_1 = 0$ ,  $E \underline{Y}_1 \underline{Y}_1' = I_p$ ,  $E \underline{Z}_1 \underline{Z}_1' = \text{diag} [\lambda_1, \dots, \lambda_p]$ ,  $E(\underline{Y}_1 \underline{Y}_1')^2 < \infty$ ,  $E(\underline{Z}_1 \underline{Z}_1')^2 < \infty$ , and that  $\lim_{n \rightarrow \infty} \frac{n}{n_1} = \theta^2 > 0$ . Then the joint distribution of  $\zeta_1^{(n)}, \dots, \zeta_p^{(n)}$  tends to that of  $\zeta_1, \dots, \zeta_p$ , where  $\zeta_{a_{h-1}+1} \geq \dots \geq \zeta_{a_h}$  are the roots of the determinant equation

$$\det (Q_{hh} - \zeta I_{\mu_h}) = 0, \quad (2.13)$$

and

$$Q_{hh} = (q_{ij}), \quad i, j = a_{h-1}+1, \dots, a_h, \quad h=1,2,\dots,H,$$

satisfying



(1)  $q_{ij} = q_{ji}$  for each  $(i,j)$ .

(2)  $\{q_{ij}, a_{h-1}+1 \leq i \leq j \leq a_h, h=1,2,\dots,H\}$  has a joint normal distribution.

(3)  $Eq_{ij} = 0$  for each  $(i,j)$ ,

$$\text{Cov}(q_{ii}, q_{jj}) = EY_{1i}^2 Y_{1j}^2 + \theta^2 \epsilon_h \epsilon_g E Z_{1i}^2 Z_{1j}^2 - \epsilon_h \epsilon_g (1 + \theta^2),$$

if  $a_{h-1} + 1 \leq i \leq a_h, a_{g-1} + 1 \leq j \leq a_g, 1 \leq h \leq g \leq H$ ,

and

$$\text{Cov}(q_{i_1 i_2}, q_{j_1 j_2}) = EY_{1i_1} Y_{1i_2} Y_{1j_1} Y_{1j_2} + \theta^2 \epsilon_h \epsilon_g E Z_{1i_1} Z_{1i_2} Z_{1j_1} Z_{1j_2},$$

if  $a_{h-1}+1 \leq i_1 \leq i_2 \leq a_h, a_{g-1}+1 \leq j_1 \leq j_2 \leq a_g, 1 \leq h \leq g \leq H$  and

at least one of the inequalities  $i_1 < i_2, j_1 < j_2$  holds.

Note that (2.13) is equivalent to (2.12) by the independence between  $U$  and  $V$ , and Theorem 2.1 holds evidently.

THEOREM 2.2. Besides the assumptions of Theorem 2.1, we assume that

$$EY_{1i}^2 Y_{1j}^2 = EY_{1i}^2 EY_{1j}^2 = 1, \quad 1 \leq i < j \leq p,$$

$$EZ_{1i}^2 Z_{1j}^2 = EZ_{1i}^2 EZ_{1j}^2 = \lambda_i \lambda_j, \quad 1 \leq i < j \leq p,$$

and

$$EY_{1i_1} Y_{1i_2} Y_{1j_1} Y_{1j_2} = EZ_{1i_1} Z_{1i_2} Z_{1j_1} Z_{1j_2} = 0$$

for all  $i_1, i_2, j_1, j_2$  satisfying that  $a_{h-1} + 1 \leq i_1 \leq i_2 \leq a_h$ ,

$a_{g-1} + 1 \leq j_1 \leq j_2 \leq a_g, 1 \leq h \leq g \leq H$  and that at least one of

$i_1 < i_2, j_1 < j_2$  holds. Then the limiting vectors  $(\tau_1, \dots, \tau_{a_1}), \dots,$

$(\tau_{a_{H-1}+1}, \dots, \tau_p)$  are independent each other.

THEOREM 2.3. Under the assumptions of Theorem 2.2 and that

$EY_{1i}^4 = 3$ ,  $EZ_{1i}^4 = 3\lambda_i^2$ ,  $i=1, \dots, p$ , the joint distribution of

$n_i^{(n)} = \sqrt{n}(\phi_i^{(n)} - \varepsilon_h) / (\sqrt{2(1+\theta^2)}\varepsilon_h)$ ,  $i=a_{h-1}+1, \dots, a_h$ ,  $h=1, 2, \dots, H$  tends to a

limiting distribution with density

$$\prod_{h=1}^H D(n_{a_{h-1}+1}, \dots, n_{a_h}),$$

where  $D(x_1, \dots, x_\mu)$  is given by

$$D(x_1, \dots, x_\mu) = 2^{-\mu/2} \left( \prod_{i=1}^{\mu} \Gamma\left(\frac{i}{2}\right) \right)^{-1} \left\{ \prod_{1 \leq i < j \leq \mu} (x_i - x_j) \right\} \exp \left\{ -\frac{1}{2} \sum_{i=1}^{\mu} x_i^2 \right\},$$

$$-\infty < x_1 \leq x_2 \leq \dots \leq x_\mu < \infty.$$

(Refer to Bai, Krishnaiah and Liang, (1984)).

To prove this theorem, it is enough to notice that  $[2(1+\theta^2)]^{-1/2} Q_{hh}$  is a (real) central Gaussian matrix of order  $\mu_h \times \mu_h$  and that  $D(x_1, \dots, x_\mu)$  is the joint density of the eigenvalues  $v_1 \geq \dots \geq v_\mu$  of a central Gaussian matrix of order  $\mu \times \mu$ . For the definition of Gaussian matrix, refer to Zhao, Krishnaiah and Bai(1987).

## 3. THE UNBALANCED CASE

In this section, we study the asymptotic joint distribution of the eigenvalues of  $\frac{N-k}{k-1} W_2 W_1^{-1}$  defined by (1.6). For simplicity, we assume that  $\alpha_j$  and  $\varepsilon_{ij}$  in (1.1) are distributed as  $p \times 1$  normal vectors for each  $(i,j)$ , and that

$$\begin{aligned} \alpha_j \text{'s are iid., } \varepsilon_{ij} \text{'s are iid., } \alpha_j \text{'s and } \varepsilon_{ij} \text{'s are} \\ \text{independent, } E(\alpha_1) = E(\varepsilon_{11}) = 0, \text{ and} \end{aligned} \quad (3.1)$$

$$E(\alpha_1 \alpha_1') = \gamma, \quad E(\varepsilon_{11} \varepsilon_{11}') = \Sigma_1 > 0.$$

Besides, we assume that

$$\frac{k}{N-k} = \theta^2 + o\left(\frac{1}{\sqrt{k}}\right), \quad \theta > 0, \text{ a constant,} \quad (3.2)$$

$$\frac{1}{k} \sum_{i=1}^k m_i^2 = m^2 + o\left(\frac{1}{\sqrt{k}}\right), \quad 1 \leq m_i \leq M, \quad i=1, \dots, k, \quad (3.3)$$

$m$  and  $M$  are constants.

Let  $\delta_1^{(k)} \geq \dots \geq \delta_p^{(k)}$ ,  $\psi_1 \geq \dots \geq \psi_p$  and  $\lambda_1^{(k)} \geq \dots \geq \lambda_p^{(k)}$  denote the eigenvalues of  $\frac{1}{k-1} W_2 (\frac{1}{N-k} W_1)^{-1}$ ,  $\gamma \Sigma_1^{-1}$  and  $\Sigma_2 \Sigma_1^{-1}$  respectively, where

$$\Sigma_2 = \Sigma_1 + \frac{N}{k} \gamma = \Sigma_1 + (1+\theta^{-2})\gamma + o\left(\frac{1}{\sqrt{k}}\right)\gamma.$$

Without loss of generality, we can assume that

$$\Sigma_1 = I_p \text{ and } \gamma = \text{diag} [\psi_1, \dots, \psi_p].$$

Write

$$\lambda_i^{(k)} = \varepsilon_h^{(k)}, \quad \psi_i = \psi_h, \quad \text{if } i = a_{h-1} + 1, \dots, a_h, \quad h=1, 2, \dots, H,$$

where  $\bar{\psi}_1 > \bar{\psi}_2 > \dots > \bar{\psi}_H \geq 0$ ,  $\varepsilon_h^{(k)} = 1 + \frac{N}{k} \bar{\psi}_h$ ,  $h=1,2,\dots,H$ .

The eigenvalues of  $\frac{1}{k-1}W_2(\frac{1}{N-k}W_1)^{-1}$  are the roots of the determinant equation

$$\det\left(\frac{1}{k-1}W_2 - \frac{\delta}{N-k}W_1\right) = 0. \quad (3.4)$$

Put  $N-k = \theta^{-2}k\beta_k^{-2}$ , then  $\beta_k = 1 + o(\frac{1}{\sqrt{k}})$  as  $k \rightarrow \infty$ .

Write

$$U_k = (u_{ij}^{(k)}) = \sqrt{N-k}\left(\frac{1}{N-k}W_1 - I_p\right),$$

$$V_k = (v_{ij}^{(k)}) = \sqrt{k}\left(\frac{1}{k-1}W_2 - \Sigma_2\right) = \sqrt{k}\left(\frac{1}{k-1}W_2 - \text{diag}[\lambda_1^{(k)}, \dots, \lambda_p^{(k)}]\right)$$

Then (3.4) becomes

$$\det\left(\frac{1}{\sqrt{k}}V_k - \frac{\theta\beta_k}{\sqrt{k}}U_k - D_k\right) = 0, \quad (3.5)$$

where

$$D_k = \begin{pmatrix} (\phi - \varepsilon_1^{(k)})I_{\nu_1} & & \\ & \ddots & \\ & & (\phi - \varepsilon_H^{(k)})I_{\nu_H} \end{pmatrix}.$$

By the central limit theorem and Lemma 2.1, we can assume that

$$\begin{aligned} U_k &\rightarrow U \triangleq (u_{ij}), \quad \text{a.s.} \\ V_k &\rightarrow V \triangleq (v_{ij}), \quad \text{a.s.} \end{aligned} \quad (3.6)$$

where  $u_{ij}$  and  $v_{ij}$  are all normal variables for all  $i,j$ , and,  $U$  and  $V$  are independent. Besides, if we split the matrices  $U_k$ ,  $V_k$ ,  $U$  and  $V$  into the following blocks

$$\begin{aligned} U_k &= (u_{hg}^{(k)}), \quad V_k = (v_{hg}^{(k)}), \quad U = (u_{hg}), \quad V = (v_{hg}), \\ h, g &= 1, 2, \dots, H, \end{aligned}$$

with  $U_{hg}^{(k)}, V_{hg}^{(k)}, U_{hg}, V_{hg}$  being of order  $\mu_h \times \mu_g$ , then  $U_{hh}, V_{hh}, h=1,2,\dots,H$ , are independent and,  $\frac{1}{\sqrt{2}} U_{hh}$  and  $(2m^2\tilde{\psi}_h^2 + 4(\theta^{-2}+1)\tilde{\psi}_h + 2)^{-1/2} V_{hh}$ ,

$h=1,2,\dots,H$ , are all central Gaussian matrices.

Let  $\zeta_i^{(k)} = \sqrt{k}(\delta_i^{(k)} - \lambda_i^{(k)})$ ,  $i=1,2,\dots,p$ . Using the argument used in the section 6, we can prove that the joint distribution of  $\zeta_1^{(k)}, \dots, \zeta_p^{(k)}$  tends to that of  $\zeta_1, \dots, \zeta_p$  as  $k \rightarrow \infty$ , where  $\zeta_{a_{h-1}+1} \geq \zeta_{a_{h-1}+2} \geq \dots \geq \zeta_{a_h}$  are the roots of

$$\det(V_{hh} - \theta \varepsilon_h U_{hh} - \zeta I_{\mu_h}) = 0, \quad h=1,2,\dots,H, \quad (3.7)$$

where

$$\varepsilon_h = 1 + (1+\theta^{-2})\tilde{\psi}_h, \quad h=1,2,\dots,H.$$

Now  $[2m^2\tilde{\psi}_h^2 + 4(\theta^{-2}+1)\tilde{\psi}_h + 2 + 2\varepsilon_h]^{-1/2}(V_{hh} - \theta \varepsilon_h U_{hh})$ ,  $h=1,2,\dots,H$ , are independent central Gaussian matrices. So we have the following

**THEOREM 3.1.** Suppose that  $\underline{\alpha}_i$  and  $\underline{\varepsilon}_{ij}$  in (1.1) are both  $p \times 1$  normal vectors for each  $(i,j)$ , and (3.1)-(3.3) are satisfied. Put

$$\eta_i^{(k)} = \sqrt{k}(\delta_i^{(k)} - \lambda_i^{(k)}) / \sqrt{2m^2\tilde{\psi}_h^2 + 4(\theta^{-2}+1)\tilde{\psi}_h + 2 + 2\varepsilon_h},$$

$$i = a_{h-1}+1, \dots, a_h, \quad h=1,2,\dots,H,$$

where  $\varepsilon_h = 1 + (1+\theta^{-2})\tilde{\psi}_h$ . Then, the joint distribution of  $\eta_1^{(k)}, \dots, \eta_p^{(k)}$  tends to a limiting distribution with density

$$\prod_{h=1}^H D(\eta_{a_{h-1}+1}, \dots, \eta_{a_h}),$$

where  $D(x_1, \dots, x_u)$  has given in Theorem 2.3.

## REFERENCES .

- [ 1 ] ANDERSON, B. M., ANDERSON, T. W. and OLKIN, I. (1986) Maximum Likelihood Estimators and Likelihood Ratio Criteria in Multivariate Components of Variance. *Ann. Statist.*, 14, No.2, 405-417.
- [ 2 ] ANDERSON, T. W. (1984). Estimating Linear Statistical Relationships (The 1982 Wald Memorial Lectures). *Ann. Statist.* 12, 1-45.
- [ 3 ] ANDERSON, T. W. (1951). The Asymptotic Distribution of Certain Characteristic Roots and Vectors. Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, Univ. of California Press, Berkeley, 103-130.
- [ 4 ] BAI, Z. D. (1984). A Note on Asymptotic Joint Distribution of the Eigenvalues of a noncentral multivariate F Matrix. Technical Report No. 84-49, Center for Multivariate Analysis, University of Pittsburgh.
- [ 5 ] BAI, Z. D., KRISHNAIAH, P. R. and LIANG, W. Q. (1984). On Asymptotic Joint Distribution of the Eigenvalues of the Noncentral MANOVA Matrix for Nonnormal Populations. Technical Report No. 84-53, Center for Multivariate Analysis, University of Pittsburgh.
- [ 6 ] BAI, Z. D., KRISHNAIAH, P. R. and ZHAO, L. C. (1986). On Multiplicities of the Eigenvalues of Component of Covariance with Applications in Signal Processing. Technical Report No. 86-37, Center for Multivariate Analysis, University of Pittsburgh.
- [ 7 ] BAI, Z. D., KRISHNAIAH, P. R. and ZHAO, L. C. (1987). Multivariate Components of Covariance Model in Unbalanced Case. To Appear in *Communications in Statistics*.
- [ 8 ] FISHER, R. A. (1939). The Sampling Distribution of some Statistics Obtained from Nonlinear Equations. *Ann. Eugen.*, 9, 238-249.
- [ 9 ] HSU, P. L. (1939). On the Distribution of Roots of Certain Determinantal Equations. *Ann. Eugen.*, 9, 250-258.
- [ 10 ] HSU, P. L. (1941). On the Limiting Distribution of Roots of a Determinantal Equations. *J. London Math. Soc.*, 16, 183-194.
- [ 11 ] RAO, C. R. (1983). Likelihood Ratio Tests for Relationships Between Two Covariance Matrices. In *Studies in Econometrics, Time Series, and Multivariate Statistics* (S. Karlin, T. Amemiya, and L. A. Goodman, eds.) 529-543. Academic, New York.
- [ 12 ] SCOTT, J. R. and SAW, J. G. (1984). A Multivariate One-way Classification Model with Random Effects. *J. Multivariate Anal.* 15, 1-12.
- [ 13 ] SKOROKHOD, A. V. (1956). Limit Theorems for Stochastic Processes. *Theory of Probability and Its Applications*, 1, 261-290.
- [ 14 ] THEOBALD, C. M. (1975). An Inequality with Application to Multivariate Analysis. *Biometrika* 62, 461-466.

- [15] ZHAO, L. C., KRISHNAIAH, P. R., AND BAI, Z. D. (1986a). On Detection of the Number of Signals in Presence of White Noise. *J. Multivariate Anal.* 20, 1-25.
- [16] ZHAO, L. C., KRISHNAIAH, P. R. AND BAI, Z. D. (1986b). On Detection of the Number of Signals when the Noise Covariance Matrix is Arbitrary. *J. Multivariate Anal.* 20, 26-49.
- [17] ZHAO, L. C., KRISHNAIAH, P. R. and BAI, Z. D. (1987). Remarks on Certain Criteria for Detection of Number of Signals. *IEEE Trans. Acoust., Speech, Signal Processing*, Vol. ASSP-35, No.2, pp.129-132.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 87-16	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On the asymptotic joint distributions of the eigenvalues of random matrices which arise under components of covariance model		5. TYPE OF REPORT & PERIOD COVERED Technical - June 1987
7. AUTHOR(s) Z. D. Bai, P. R. Krishnaiah and L. C. Zhao		6. PERFORMING ORG. REPORT NUMBER 87-16
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis 515 Thackeray Hall University of Pittsburgh, Pittsburgh, PA 15260		8. CONTRACT OR GRANT NUMBER(s) F49620-85-C-0008
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Rolling Air Force Base, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1987
		13. NUMBER OF PAGES 16
		15. SECURITY CLASS. (of this report) Unclassified
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Component of covariance model, eigenstructure analysis, limiting distribution, random matrix.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  In this paper, the authors derived asymptotic joint distributions of the eigenvalues of some random matrices which arise under components of covariance model.		



END

DATE

FILMED

JAN

1988